



BOTOX® Complete offers support at every step—for everyone

If you're new to BOTOX®, enroll for

- ✓ Tools to help you start and stay on track with your treatment
- ✓ Potential opportunities to save
- ✓ BOTOX® information

Enroll in BOTOX® Complete



Visit **BOTOX.com/account**
or text **ENROLL** to 27747*

SAFETY CONSIDERATIONS

Effects of BOTOX® may spread hours to weeks after injection, causing serious symptoms. Alert your doctor right away, as difficulty swallowing, speaking, and breathing can be signs of a life-threatening condition. Patients with these conditions before injection are at highest risk. In some cases, the effect of botulinum toxin may spread to other areas of the body, causing serious symptoms, such as muscle weakness, vision changes, eye problems, changes or loss of voice, and loss of bladder control. Side effects may include: allergic reactions; neck and injection-site pain; tiredness; headache; eye problems; drooping eyebrows; upper respiratory tract infection; dry mouth; in adults being treated for urinary incontinence: UTI, inability to empty your bladder on your own, and painful urination; and in pediatric patients being treated for urinary incontinence: UTI, inability to empty your bladder on your own, and bacteria, white blood cells, and blood in the urine. Don't receive BOTOX® if there's an infection at the planned injection site, you are allergic to any of the ingredients in BOTOX®, you have had an allergic reaction to any other botulinum toxin product, or you cannot empty your bladder on your own (in those being treated for urinary incontinence who are not routinely catheterizing).

*Msg&data rates may apply. You are not required to consent as a condition of receiving goods or services. You can reply HELP for help. You can reply STOP to opt out at any time. By texting ENROLL to 27747, you agree to AbbVie's SMS Terms and Conditions at www.botoxsms.com and Privacy Notice at <http://abbvie.com/corpprivacy>.

Please see **Important Safety Information** on pages 3–4.

Please see full **Prescribing Information**, including Boxed Warning and Medication Guide, or visit https://www.rxabbvie.com/pdf/botox_pi.pdf.

Stay connected throughout your prescribed treatment

BOTOX® Complete helps you navigate challenges as they arise and find answers when you need them. When you enroll, you can expect:



Regular support on your personalized dashboard



Help getting started with treatment



Savings opportunities for eligible, commercially insured patients



Treatment reminders to help you stay on track

Enroll in BOTOX® Complete



Visit **BOTOX.com/account** or text **ENROLL** to 27747*

You may pay as little as \$0†

Eligible, commercially insured patients could pay as little as \$0† for their BOTOX® treatment and receive reimbursement for out-of-pocket costs‡



Submit a reimbursement claim

When you submit a claim for reimbursement on your BOTOX® Complete dashboard, you could save on costs related to both BOTOX® medicine and the injection procedure.‡




Use your Specialty Pharmacy Savings Card

If you fill your BOTOX® prescription with a Specialty Pharmacy, they will automatically apply the eligible benefit from your Savings Card toward your out-of-pocket costs.‡



Not actual Savings Card.



Enroll in BOTOX® Complete to see if you're eligible

*Msg&data rates may apply. You are not required to consent as a condition of receiving goods or services. You can reply HELP for help. You can reply STOP to opt out at any time. By texting ENROLL to 27747, you agree to AbbVie's SMS Terms and Conditions at www.botoxsm.com and Privacy Notice at <http://abbvie.com/privacy>.

†Eligibility: Available to patients with commercial insurance coverage for BOTOX® (onabotulinumtoxinA) who meet eligibility criteria. This co-pay assistance program is not available to patients receiving prescription reimbursement under any federal, state, or government-funded insurance programs (for example, Medicare [including Part D], Medicare Advantage, Medigap, Medicaid, TRICARE, Department of Defense, or Veterans Affairs programs) or where prohibited by law. Offer subject to change or termination without notice. Restrictions, including monthly maximums, may apply. This is not health insurance. **For full Terms and Conditions**, visit <https://www.botox.com/completeterms> or call 1-800-44-BOTOX (800-442-6869) for additional information. To learn about AbbVie's privacy practices and your privacy choices, visit <https://abbvie.com/privacy>.

‡For residents of some states, offer applies only to the cost of BOTOX® and not to any related medical service(s).

Please see **Important Safety Information** on pages 3–4.

Please see full **Prescribing Information**, including Boxed Warning and Medication Guide, or visit https://www.rxabbvie.com/pdf/botox_pi.pdf.

INDICATIONS

BOTOX® (onabotulinumtoxinA) is a prescription medicine that is injected into muscles and used:

- To treat overactive bladder symptoms such as a strong need to urinate with leaking or wetting accidents (urge urinary incontinence), a strong need to urinate right away (urgency), and urinating often (frequency) in adults 18 years and older when another type of medicine (anticholinergic) does not work well enough or cannot be taken
- To treat leakage of urine (incontinence) in adults 18 years and older with overactive bladder due to a neurologic disease when another type of medicine (anticholinergic) does not work well enough or cannot be taken
- To treat overactive bladder due to a neurologic disease in children 5 years of age and older when another type of medicine (anticholinergic) does not work well enough or cannot be taken
- To prevent headaches in adults with chronic migraine who have 15 or more days each month with headache lasting 4 or more hours each day in people 18 years and older
- To treat increased muscle stiffness in people 2 years of age and older with spasticity
- To treat the abnormal head position and neck pain that happens with cervical dystonia (CD) in people 16 years and older
- To treat certain types of eye muscle problems (strabismus) or abnormal spasm of the eyelids (blepharospasm) in people 12 years of age and older

BOTOX is also injected into the skin to treat the symptoms of severe underarm sweating (severe primary axillary hyperhidrosis) when medicines used on the skin (topical) do not work well enough in people 18 years and older.

It is not known whether BOTOX is safe and effective to prevent headaches in patients with migraine who have 14 or fewer headache days each month (episodic migraine).

BOTOX has not been shown to help people perform task-specific functions with their upper limbs or increase movement in joints that are permanently fixed in position by stiff muscles.

It is not known whether BOTOX is safe and effective for severe sweating anywhere other than your armpits.

IMPORTANT SAFETY INFORMATION

BOTOX may cause serious side effects that can be life threatening. Get medical help right away if you have any of these problems any time (hours to weeks) after injection of BOTOX:

- **Problems swallowing, speaking, or breathing**, due to weakening of associated muscles, can be severe and result in loss of life. You are at the highest risk if these problems are preexisting before injection. Swallowing problems may last for several months.
- **Spread of toxin effects**. The effect of botulinum toxin may affect areas away from the injection site and cause serious symptoms, including loss of strength and all-over muscle weakness; double vision; blurred vision; drooping eyelids; hoarseness or change or loss of voice; trouble saying words clearly; loss of bladder control; trouble breathing; and trouble swallowing.

There has not been a confirmed serious case of spread of toxin effect away from the injection site when BOTOX has been used at the recommended dose to treat chronic migraine, severe underarm sweating, blepharospasm, or strabismus.

BOTOX may cause loss of strength or general muscle weakness, vision problems, or dizziness within hours to weeks of receiving BOTOX. **If this happens, do not drive a car, operate machinery, or do other dangerous activities.**

Do not receive BOTOX if you are allergic to any of the ingredients in BOTOX (see Medication Guide for ingredients); had an allergic reaction to any other botulinum toxin product such as Myobloc® (rimabotulinumtoxinB), Dysport® (abobotulinumtoxinA), Xeomin® (incobotulinumtoxinA), Jeuveau® (prabotulinumtoxinA-xvfs), Daxxify® (daxibotulinumtoxinA-lanm), or Letybo® (letibotulinumtoxinA-wlbq) (this may not be a complete list of all botulinum toxin products); have a skin infection at the planned injection site.

Do not receive BOTOX for the treatment of urinary incontinence if you have a urinary tract infection (UTI) or cannot empty your bladder on your own and are not routinely catheterizing. Due to the risk of urinary retention (difficulty fully emptying the bladder), only patients who are willing and able to initiate catheterization posttreatment, if required, should be considered for treatment.

Patients treated for overactive bladder: In clinical trials, 36 of the 552 patients had to self-catheterize for urinary retention following treatment with BOTOX compared to 2 of the 542 treated with placebo. The median duration of postinjection catheterization for these patients treated with BOTOX 100 Units (n = 36) was 63 days (minimum 1 day to maximum 214 days), as compared to a median duration of 11 days (minimum 3 days to maximum 18 days) for patients receiving placebo (n = 2). Patients with diabetes mellitus treated with BOTOX were more likely to develop urinary retention than nondiabetics.

Adult patients treated for overactive bladder due to a neurologic disease: In clinical trials, 30.6% of adult patients (33/108) who were not using clean intermittent catheterization (CIC) prior to injection required catheterization for urinary retention following treatment with BOTOX 200 Units, as compared to 6.7% of patients (7/104) treated with placebo. The median duration of postinjection catheterization for these patients treated with BOTOX 200 Units (n = 33) was 289 days (minimum 1 day to maximum 530 days), as compared to a median duration of 358 days (minimum 2 days to maximum 379 days) for patients receiving placebo (n = 7).

Please see **Important Safety Information** on pages 3–4.

Please see full **Prescribing Information**, including **Boxed Warning** and **Medication Guide**, or visit https://www.rxabbvie.com/pdf/botox_pi.pdf.


onabotulinumtoxinA_{injection}

IMPORTANT SAFETY INFORMATION (continued)

Among adult patients not using CIC at baseline, those with multiple sclerosis were more likely to require CIC postinjection than those with spinal cord injury.

The dose of BOTOX is not the same as, or comparable to, another botulinum toxin product.

Serious and/or immediate allergic reactions have been reported, including itching; rash; red, itchy welts; wheezing; asthma symptoms; dizziness; or feeling faint. Get medical help right away if you experience symptoms; further injection of BOTOX should be discontinued.

Tell your doctor about all your muscle or nerve conditions, such as ALS or Lou Gehrig's disease, myasthenia gravis, or Lambert-Eaton syndrome, as you may be at increased risk of serious side effects, including difficulty swallowing and difficulty breathing from typical doses of BOTOX.

Tell your doctor if you have any breathing-related problems. Your doctor may monitor you for breathing problems during treatment with BOTOX for spasticity or for detrusor overactivity associated with a neurologic condition. The risk of developing lung disease in patients with reduced lung function is increased in patients receiving BOTOX.

Cornea problems have been reported. Cornea (surface of the eye) problems have been reported in some people receiving BOTOX for their blepharospasm, especially in people with certain nerve disorders. BOTOX may cause the eyelids to blink less, which could lead to the surface of the eye being exposed to air more than is usual. Tell your doctor if you experience any problems with your eyes while receiving BOTOX. Your doctor may treat your eyes with drops, ointments, contact lenses, or with an eye patch.

Bleeding behind the eye has been reported. Bleeding behind the eyeball has been reported in some people receiving BOTOX for their strabismus. Tell your doctor if you notice any new visual problems while receiving BOTOX.

Bronchitis and upper respiratory tract infections (common colds) have been reported. Bronchitis was reported more frequently in adults receiving BOTOX for upper limb spasticity. Upper respiratory infections were also reported more frequently in adults with prior breathing-related problems with spasticity. In pediatric patients treated with BOTOX for upper limb spasticity, upper respiratory tract infections were reported more frequently. In pediatric patients treated with BOTOX for lower limb spasticity, upper respiratory tract infections were not reported more frequently than placebo.

Autonomic dysreflexia in patients treated for overactive bladder due to a neurologic disease. Autonomic dysreflexia associated with intradetrusor injections of BOTOX could occur in patients treated for detrusor overactivity associated with a neurologic condition and may require prompt medical therapy. In clinical trials, the incidence of autonomic dysreflexia was greater in adult patients treated with BOTOX 200 Units compared with placebo (1.5% versus 0.4%, respectively).

Tell your doctor about all your medical conditions, including if you have or have had bleeding problems; have plans to have surgery; had surgery on your face; have weakness of forehead muscles, trouble raising your eyebrows, drooping eyelids, and any other abnormal facial change; have symptoms of a UTI and are being treated for urinary incontinence (symptoms of a UTI may include pain or burning with urination, frequent urination, or fever); have problems emptying your bladder on your own and are being treated for urinary incontinence; are pregnant or plan to become pregnant (it is not known if BOTOX can harm your unborn baby); are breastfeeding or plan to (it is not known if BOTOX passes into breast milk).

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using BOTOX with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you have received BOTOX in the past.**

Tell your doctor if you received any other botulinum toxin product in the last 4 months; have received injections of botulinum toxin such as Myobloc®, Dysport®, Xeomin®, Jeuveau®, Daxxify®, or Letybo® in the past (this may not be a complete list of all botulinum toxin products; tell your doctor exactly which product you received); have recently received an antibiotic by injection; take muscle relaxants; take an allergy or cold medicine; take a sleep medicine; take aspirin-like products or blood thinners.

Other side effects of BOTOX include dry mouth; discomfort or pain at the injection site; tiredness; headache; neck pain; eye problems such as double vision, blurred vision, decreased eyesight, drooping eyelids, swelling of your eyelids, and dry eyes; drooping eyebrows; and upper respiratory tract infection. In adults being treated for urinary incontinence, other side effects include UTI and painful urination. In children being treated for urinary incontinence, other side effects include UTI; bacteria, white blood cells, and blood in the urine. In patients being treated for urinary incontinence, another side effect includes the inability to empty your bladder on your own. If you have difficulty fully emptying your bladder on your own after receiving BOTOX, you may need to use disposable self-catheters to empty your bladder up to a few times each day until your bladder is able to start emptying again.

For more information, refer to the Medication Guide or talk with your doctor.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/PatientAccessSupport to learn more.

abbvie

© 2026 AbbVie. All rights reserved.

BOTOX® and its design are registered trademarks of Allergan, Inc., an AbbVie company. All other trademarks are the property of their respective owners.

US-BTX-250353


onabotulinumtoxinA_{injection}